

E. Ballico

Symmetric tensor rank over an infinite field

Received: 2 March 2013 / Accepted: 28 May 2013 / Published online: 13 June 2013
 © The Author(s) 2013. This article is published with open access at Springerlink.com

Abstract Here we prove two upper bounds (one for bivariate polynomials, one for multivariate ones) for the symmetric tensor rank with respect to an infinite field with characteristic $\neq 2$.

Mathematics Subject Classification 14N05 · 14Q05 · 15A69

المخلص

نثبت هنا وجود حدين علويين (أحدهما لكثيرات الحدود ثنائية المتغير والآخر لكثيرات الحدود متعددة المتغيرات) لرتبة المؤثر المتناظر بالنسبة لحقل لانهاضي مميزه $\neq 2$.

1 Introduction

There is a huge and fast growing literature on the rank of tensors and on the symmetric tensor rank of symmetric tensors [4, 6, 9, 10, 12, 14–16]. There are real world applications and motivations (e.g. signal processing) behind these papers. Most of the papers are over \mathbb{C} (or over an algebraically closed field), but real tensors and real polynomials are also quite studied [13, 16]. Here, we work over an arbitrary infinite field and extend to the non-algebraically closed case, a known upper bound for the symmetric tensor rank.

Let K be a field and \bar{K} its algebraic closure. For any $A \subseteq \mathbb{P}^r(K)$ let $\langle A \rangle_K$ denote the K -linear span of A , i.e., the minimal K -linear subspace of $\mathbb{P}^r(K)$ containing A . Set $\langle A \rangle := \langle A \rangle_{\bar{K}}$. Notice that $\langle A \rangle_K = \langle A \rangle \cap \mathbb{P}^r(K)$ for any set $A \subseteq \mathbb{P}^r(K)$. Fix an integer $r > 0$ and a set $Y \subset \mathbb{P}^r(K)$. For each $P \in \mathbb{P}^r(K)$ the rank $r_Y(P)$ of P with respect to Y is the minimal cardinality of a set $S \subseteq Y$ such that $P \in \langle S \rangle_K$ (or, equivalently, $P \in \langle S \rangle$), with the convention $r_Y(P) = +\infty$ if $P \notin \langle Y \rangle_K$. Take a projective variety $X \subseteq \mathbb{P}^r$ defined over K . We have two sets $X(K) \subseteq \mathbb{P}^r(K)$ and $X(\bar{K}) \subseteq \mathbb{P}^r(\bar{K})$. Obviously $r_{X(K)}(P) \geq r_{X(\bar{K})}(P)$ for any $P \in \mathbb{P}^r(K)$. Quite often the latter inequality is a strict inequality (even when $K = \mathbb{R}$) [13]. Recently we realized that in the case of the symmetric tensor rank (see below for its definition) many known upper bounds for $r_{X(\bar{K})}(P)$ hold over K assuming that K is infinite. We first prove the following result.

Theorem 1.1 Fix an integer $d \geq 2$. Let K be a field such that $\text{char}(K) \neq 2$ and either K is infinite or $\sharp(K) \geq 2d - 2$. Let $C \subset \mathbb{P}^d$ be a degree d linearly normal embedding of \mathbb{P}^1 defined over K (equivalently, C is a rational normal curve of \mathbb{P}^d defined over K and with $C(K) \neq \emptyset$). Then

- (a) $r_{C(K)}(P) \leq d$ for all $P \in \mathbb{P}^d(K)$.
- (b) For any $A \subset C(K)$ such that $\sharp(A) = d - 2$ there is $S \subset C(K)$ such that $A \subset S$, $\sharp(S) = d$ and $P \in \langle S \rangle_K$.

E. Ballico (✉)
 Department of Mathematics, University of Trento, 38123 Povo (TN), Italy
 E-mail: ballico@science.unitn.it



- (c) Assume that K is infinite. For any finite $B \subset C(K)$ and any $A \subset C(K)$ such that $\sharp(A) = d - 2$ and $A \cap B = \emptyset$ there is $S \subset C(K)$ such that $A \subset S$, $B \cap S = \emptyset$, $\sharp(S) = d$ and $P \in \langle S \rangle_K$.

Theorem 1.1 concerns the one-dimensional case (i.e., the bivariate case) of the following set-up.

Fix an infinite field K and positive integers m, d . Set $N := \binom{m+d}{m} - 1$. For any field $L \supseteq K$ let $v_{d,L} : \mathbb{P}^m(L) \rightarrow \mathbb{P}^N(L)$ denote the order d Veronese embedding of $\mathbb{P}^m(L)$ obtained using the L -vector space of all degree d homogeneous polynomials in $m + 1$ variables. Set $X_{m,d}(L) := v_{d,L}(\mathbb{P}^m(L))$. The set $X_{m,d}(L)$ is the set of L -points of the order d Veronese embedding $X_{m,d}$ of \mathbb{P}^m . Notice that $v_{d,K} = v_{d,L}|_{\mathbb{P}^m(K)}$. Since L is infinite, $\mathbb{P}^m(L)$ is Zariski dense in $\mathbb{P}^m(\bar{L})$. Since L is infinite, for every $f \in L[x_0, \dots, x_m] \setminus \{0\}$ there is $(b_0, \dots, b_m) \in L^{m+1}$ such that $f(b_0, \dots, b_m) \neq 0$. Hence the set $X_{m,d}(L)$ spans $\mathbb{P}^N(L)$. Hence for each $P \in \mathbb{P}^N(L)$, $sr_L(P) := r_{X_{m,d}(L)}(P)$ is a positive integer, often called the L -symmetric tensor rank of P . Now assume $P \in \mathbb{P}^N(K)$. Since $X_{m,d}(L) \supseteq X_{m,d}(K)$, we have $sr_L(P) \leq sr_K(P)$. Hence $sr_K(P) \geq sr_{\bar{K}}(P)$. The integer $sr(P) := sr_{m,d,\bar{K}}(P)$ is the usual symmetric tensor rank of P [14]. Using Theorem 1.1, we are able to prove the following extension of Ballico [5, Theorem 1], to a non-algebraically closed field.

Theorem 1.2 Fix integers m, d such that $m \geq 5$ and $d \geq 3$. Let K be an infinite field such that $\text{char}(K) \neq 2$. For any $P \in \mathbb{P}^N(K)$ we have

$$sr_K(P) \leq d \cdot \lceil \binom{m+d}{m} / (m+1) \rceil.$$

The proof of the case $m \geq 2$ of Theorem 1.2 is only a minor modification of the proof of [5, Theorem 1]. For multivariate polynomials, Białynicki-Birula and Schinzel [7,8] proved a very good upper bound for the symmetric tensor rank. Assume $\text{char}(\bar{K}) = 0$. Let $X(\bar{K}) \subset \mathbb{P}^n(\bar{K})$ be an integral and non-degenerate variety defined over \bar{K} . Landsberg and Teitler [15, Proposition 5.1] proved that $r_{X(\bar{K})}(P) \leq n - \dim(X) + 1$ for all $P \in \mathbb{P}^n(\bar{K})$. This bound is sharp for bivariate polynomials (i.e., if $X(\bar{K})$ a rational normal curve) and for several other curves.

2 The proofs

Proof of Theorem 1.1. We first check part (b) and hence part (a).

Since $C(K) \neq \emptyset$, either $C(K)$ is infinite (case K infinite) or $\sharp(C(K)) = \sharp(K) + 1$ (case K finite). Fix $P \in \mathbb{P}^d(K)$ and $A \subset C(K)$ such that $\sharp(A) = d - 2$. We have $\sharp(C(K) \setminus A) \geq d + 1$. Since C is a rational normal curve, we have $\dim(\langle Z \rangle) = \deg(Z) - 1$ for every zero-dimensional scheme $Z \subset C(\bar{K})$ with $\deg(Z) \leq d + 1$. Hence $\dim(\langle A \rangle_K) = d - 3$ and $\langle A \rangle \cap C(\bar{K}) = A$ as schemes. If $P \in \langle A \rangle$, then we are done. Hence we may assume $\dim(\langle \{P\} \cup A \rangle) = d - 2$. Notice that $\langle \{P\} \cup A \rangle \cap C(\bar{K})$ has degree $\leq d - 1$ and it is defined over K . Let $\ell : \mathbb{P}^d(\bar{K}) \setminus \langle A \rangle_{\bar{K}} \rightarrow \mathbb{P}^2(\bar{K})$ denote the linear projection from $\langle A \rangle_{\bar{K}}$. Since $\langle A \rangle \cap C(\bar{K}) = A$, the restriction of ℓ induces a morphism $\eta : C(\bar{K}) \setminus A \rightarrow \mathbb{P}^2(\bar{K})$. Let $Y(\bar{K})$ be the closure of $\eta(C(\bar{K}) \setminus A)$ in $\mathbb{P}^2(\bar{K})$. Since $\dim(\langle Z \rangle) = \deg(Z) - 1$ for every zero-dimensional scheme $Z \subset C(\bar{K})$ with $\deg(Z) \leq d + 1$, $Y(\bar{K})$ is a smooth conic and η is injective. Since ℓ is defined over K , η is defined over K . For each $Q \in A$ set $A_Q := (A \setminus \{Q\}) \cup 2Q$ seen as a degree $d - 1$ effective divisor of Q . Since $\dim(\langle Z \rangle) = \deg(Z) - 1$ for every zero-dimensional scheme $Z \subset C(\bar{K})$ with $\deg(Z) \leq d$, $Y(\bar{K}) \setminus \eta(C(\bar{K}) \setminus A)$ is formed by $d - 2$ points, each of them defined over K and each of them being one of the points $\ell(\langle A_Q \rangle \setminus \langle A \rangle)$, $Q \in A$. Hence $\ell(P) \notin Y(\bar{K}) \setminus \eta(C(\bar{K}) \setminus A)$. Since $\text{char}(K) \neq 2$, for any smooth conic $E \subset \mathbb{P}^2(\bar{K})$ and any $O \in \mathbb{P}^2(\bar{K})$, there are exactly two tangent lines of E containing O . Since $\text{char}(K) \neq 2$ and $Y(\bar{K})$ is a smooth conic, there are at most two tangent lines to $Y(\bar{K})$ passing through $\ell(P)$. Since $\eta(C(K) \setminus A)$ has at least cardinality 3, there is a line $T \subset \mathbb{P}^2(\bar{K})$ containing $\ell(P)$, a point $B \in \eta(C(K) \setminus A)$ and not tangent to $Y(\bar{K})$. Since $\ell(P)$ and B are defined over K , the line T is defined over K . Hence the other point, B' , of $T \cap Y(\bar{K})$ is defined over K . Notice that B' is uniquely determined by P and B . Since $\sharp(\eta(C(K) \setminus A)) \geq d + 1$, we may find B with the additional condition that $B' \notin Y(\bar{K}) \setminus \eta(C(\bar{K}) \setminus A)$. Hence there are $O \in C(K) \setminus A$ and $O' \in C(K) \setminus A$ such that $\ell(O) = B$ and $B' = \ell(O')$. Since $\ell(P) \in T(K) = \langle \{B, B'\} \rangle_K$, we have $P \in \langle A \cup \{O, O'\} \rangle_K$. Take $S := A \cup \{O, O'\}$.

Now assume that K is infinite and take a finite $B \subset C(K)$ such that $B \cap A = \emptyset$. Since $A \cap B = \emptyset$, the set $\ell(B)$ is well defined. Since $\ell(B)$ is finite, we may find T as above with the additional condition $T \cap \ell(B) = \emptyset$. \square



Proof of Theorem 1.2. Set $\alpha := \lceil \binom{m+d}{m} / (m+1) \rceil$. Fix $P \in \mathbb{P}^N(K)$.

First assume $\text{char}(K) = 0$. For each $U \in X_{n,d}(\overline{K})$ let $T_U X_{m,d}$ be the tangent space of $X_{m,d}(\overline{K})$ at U seen as an m -dimensional \overline{K} -linear subspace of $\mathbb{P}^N(\overline{K})$. For any $U \in \mathbb{P}^N$ set $T_{U,K} X_{m,d} := T_U X_{m,d} \cap \mathbb{P}^N(K)$. Since the morphism $v_{d,\overline{K}}$ is defined over K , $T_{U,K} X_{m,d}$ is an m -dimensional K -linear subspace. Since $m \geq 5$ and $d \geq 3$, a theorem of J. Alexander and A. Hirschowitz [2,3] says that α is the first positive integer t such that $\langle \cup_{Q \in B} T_{v_d(Q)} X_{m,d} \rangle = \mathbb{P}^N(\overline{K})$ for a general $B \subset \mathbb{P}^m(\overline{K})$ with cardinality t (Terracini's lemma [1], part 2 of Corollary 1.11). In positive characteristic only one implication holds in Terracini's lemma [1, part 1 of Corollary 1.11]. This is the version of Alexander–Hirschowitz theorem proved in arbitrary characteristic by Chandler [11, Theorem 1]. Hence in arbitrary characteristic $\langle \cup_{Q \in B} T_{v_d(Q)} X_{m,d} \rangle = \mathbb{P}^N(\overline{K})$ for a general $B \subset \mathbb{P}^m(\overline{K})$ with cardinality α . Since K is infinite, $\mathbb{P}^m(K)$ is Zariski dense in $\mathbb{P}^m(\overline{K})$. Hence there is $B \subset \mathbb{P}^m(K)$ such that $\langle \cup_{Q \in B} T_{v_d(Q)} X_{m,d} \rangle = \mathbb{P}^N(\overline{K})$. Hence $\langle \cup_{Q \in B} T_{v_d(Q),K} X_{m,d} \rangle = \mathbb{P}^N(K)$. Hence there are $O_b \in T_{v_d(B),K} X_{m,d}$, $b \in B$, such that $P \in \langle \cup_{b \in B} O_b \rangle$. Fix $b \in B$. Let $2b$ be the closed subscheme of $\mathbb{P}^m(\overline{K})$ with $(\mathcal{I}_b)^2$ as its ideal sheaf. The scheme $2b$ is a degree $m+1$ zero-dimensional scheme defined over K . Hence it is defined the K -linear space $\langle v_d(2b) \rangle_K$. Since $v_{d,\overline{K}}$ is an embedding, we have $\dim(\langle v_d(2b) \rangle_K) = m+1$. We have $\langle v_d(2b) \rangle_K = T_{v_d(b),K} X_{m,d}$. Let $J \subset \mathbb{P}^N(\overline{K})$ be the line spanned by $v_{d,K}(b)$ and by O_b . Since $\{v_{d,K}(b), O_b\} \subset \mathbb{P}^N(K)$, the line J is defined over K . Hence the scheme $Z := v_{d,K}(2b) \cap J$ is defined over K . We have $\deg(Z) = 2$. Hence there is a degree 2 zero-dimensional scheme $W \subset 2b$ such that $O_b \in \langle v_d(W) \rangle$. Notice that $\langle v_d(W) \rangle$ is the line J spanned by $v_d(b)$ and by O_b and that $\deg(Z) = \deg(W) = 2$. Since b and O_b are defined over K , the line $\langle v_d(Z_b) \rangle$ is defined over K . Hence $\langle v_d(W) \rangle_K$ is a well-defined K -line. Set $D := \langle W \rangle \subset \mathbb{P}^m(\overline{K})$. D is a line defined over K . Hence $C := v_{d,\overline{K}}(D)$ is a rational normal curve defined over K . Since $b \in B$, we have $C(K) \neq \emptyset$. Since $O_b \in \mathbb{P}^N(K)$, we have $O_b \in \langle v_d(W) \rangle \cap \mathbb{P}^N(K) = \langle v_d(W) \rangle_K$. Since $O_b \in \langle v_d(D_b) \rangle_K$, there is $S_b \subset \mathbb{P}^m(K)$ such that $\#(S_b) \leq d$ and $O_b \in \langle v_d(S_b) \rangle_K$ (Theorem 1.1; only here we use the assumption $\text{char}(K) \neq 2$). Hence the set $\cup_{b \in B} S_b$ proves the inequality $sr_K(P) \leq d\alpha$. \square

Question 1 Fix an integer $d \geq 2$ and a prime p . If $p = 2$, then assume $d \geq 3$. Which is the minimal p -power $\alpha(d, p)$ such that Theorem 1.1 is true for the field \mathbb{F}_q for all p -powers $q \geq \alpha(d, p)$?

Acknowledgments The author was partially supported by MIUR and GNSAGA of INdAM (Italy).

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

1. Ådlandtsvik, B.: Joins and higher secant varieties. *Math. Scand.* **61**, 213–222 (1987)
2. Alexander, J.; Hirschowitz, A.: La méthode d'Horace éclatée: application à l'interpolation en degré quatre. *Invent. Math.* **107**, 585–602 (1992)
3. Alexander, J.; Hirschowitz, A.: Polynomial interpolation in several variables. *J. Algebraic Geom.* **4**(2), 201–222 (1995)
4. Ballico, E.; Bernardi, A.: Decomposition of homogeneous polynomials with low rank. *Math. Z.* **271**(3–4), 1141–1149 (2012). doi:10.1007/s00209-011-0907-6
5. Ballico, E.: An upper bound for the symmetric tensor rank of a low degree polynomial in a large number of variables. *ISRN-Geometry*, vol. 2013. Article ID 715907. doi:10.1155/2013/715907
6. Bernardi, A.; Gimigliano, A.; Idà, M.: Computing symmetric rank for symmetric tensors. *J. Symb. Comput.* **46**(1), 34–53 (2011)
7. Białynicki-Birula, A.; Schinzel, A.: Representations of multivariate polynomials by sums of univariate polynomials in linear forms. *Colloq. Math.* **112**(2), 201–233 (2008)
8. Białynicki-Birula, A.; Schinzel, A.: Corrigendum to “Representatons of multivariate polynomials by sums of univariate polynomials in linear forms”. *Colloq. Math.* **125**(1), 139 (2011)
9. Brachat, J.; Comon, P.; Mourrain, B.; Tsigras, E.P.: Symmetric tensor decomposition. *Linear Algebra Appl.* **433**(11–12), 1851–1872 (2010)
10. Buczyński, J.; Landsberg, J.M.: Rank of tensors and a generalization of secant varieties. *Linear Algebra Appl.* **438**(2), 668–689 (2013)
11. Chandler, K.: A brief proof of a maximal rank theorem for generic double points in projective space. *Trans. Am. Math. Soc.* **353**(5), 1907–1920 (2000)
12. Comon, P.; Golub, G.H.; Lim, L.-H.; Mourrain, B.: Symmetric tensors and symmetric tensor rank. *SIAM J. Matrix Anal.* **30**(3), 1254–1279 (2008)
13. Comon, P.; Ottaviani, G.: On the typical rank of real binary forms. *Linear Multilinear Algebra* **60**(6), 657–667 (2012)



14. Landsberg, J.M.: Tensors: Geometry and Applications. Graduate Studies in Mathematics, vol. 128. American Mathematical Society, Providence (2012)
15. Landsberg, J.M.; Teitler, Z.: On the ranks and border ranks of symmetric tensors. *Found. Comput. Math.* **10**(3), 339–366 (2010)
16. Lim, L.H.; de Silva, V.: Tensor rank and the ill-posedness of the best low-rank approximation problem. *SIAM J. Matrix Anal. Appl.* **30**(3), 1084–1127 (2008)

